3:40am brought back to sit in a chair, just need medication or general advice/recommendation for this issue - extreme anal expulsive n/ 1" + anal prolapse - possible Rectal Cancer? (what creates the stretching feeling? Many disease/disorders or just a few [FASTEST RESPONSE] -- GET TO THE ROOT - POSSIBILITIES - BE TRANSPARENT - OPEN FOR PEER REVIEW // sentence whole class (H.S.) & college/towns/visited locations & peoples who prolonged & increased prolapse of anus;

- You guys dont know where this shit came from?"
- Spencer!" "coming" (1/2 hrs later while team talks in excitement "cl will find a horse!")

Q's
- ☐ Do cl have Rectal Cancer
- ☐ Why do cl have to sit & wait on this "team"?

Blood taken @ 4:50AM
no food, for 6 hrs (bag locked away)

- Did your Medical Degree make you feel smarter or care more? work harder?
- ☐ R.O.I. for Vancouver, BC 2 injections

ii

(5/1/2018)
- ☒ emailed end the schism@gmail.com "my chart" health insurance for Washington
- ☒ Dont see the writings as personal BUT Medical
- 7:15, next bus 9am: Blaine bus 75 // ☒ Black jeep
- Steve-o saves the day, while gazing @ a star

// PTSD root function of most all mental disorders
- ☐ Monday DHS office ☐ pay check (cash assistance) ☐ Food stamp ☐ I.D.

// ☐ Run (lets try tomorrow morn, over to BMX n back)

//

5/1 iii

The Truthfulness of:
- have cl listened to tool "allot" a good amount
- ☒ Schism by Tool
- ☒ metallica ☒ clear water
- ☒ ODESSA revival
- // The fucked up thing about the Human Spirit - to giggle @ extreme trauma (nightmare - gallows humor -
- Q: = Why do we laugh @ our own futility? inability to actually effect/change

5/1 ~~11~~ 30 - B iv

# PREVENTING CHILD WINDOW FALLS

With warmer weather, windows open up and the risk for window falls increases. According to a report by Safe Kids Worldwide, an average of eight children age 5 and younger die and more than 3,300 are injured each year from falling out of windows. Falls are the most common unintentional injury requiring hospitalization for children.

Screens are not a substitute for window guards or stops, as they're not designed to prevent children from falling out windows. Children under age seven are at particular risk, and caregivers are often present at the time of the fall, so supervision alone is not enough to prevent a window fall. (Safekids.org)

Safe Kids surveyed parents about home window safety. Seventy percent of parents surveyed say they have never used window guards or stops that prevent windows from opening (Safekids.org).

Safe Kids Survey Question: Do you use window guards or stops that prevent windows from opening far enough for a child to fall out, or have you used them in the past?


Yes, I have used window guards on all windows above the first floor
Yes, I have used window guards on some windows
I'm not sure
Yes, I have used window guards on all windows
No, I've never used window guards
4%
5%
7%
14%
70%


0 in.
1 in.
2 in.
3 in.
4 in.
Screens keep bugs out, not kids in.
Install window stops so that windows do not open more than 4 inches.
Actively supervise kids around open windows.
Move furniture away from windows to prevent kids from climbing.

Learn more about securing your home against child window falls with these **Window Safety Tips**:

**Don't rely on insect screens to protect kids from falling from open windows; they are designed to keep bugs out, not to keep children in.**

Install window stops, which limit window openings to 4 inches or less, to child-accessible windows. Make sure to purchase one with a quick-release feature so adults can open the window to escape during a fire or emergency.

- Install an operable window guard if you must open a window within a child's reach. Choose window guards that can be removed by an adult in an emergency.
- Don't place furniture such as chairs, sofas and tables near windows. Remove accessibility to open windows

for ventilation, use windows
e reach of children.

y near windows, balconies,
't allow children to jump on

To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email doh.information@doh.wa.gov.

Washington State Department of HEALTH

locals, fearful in disguise, ilk and fry, like nasty lil flies (no, d dont love all my babies, see: salt lake city mother of mayne (wild & trapped) /// my neighbors → /baby nightmares ///

The Greatest Savior ran Nike a Shining Light in Dark times, gazing up at a Savior - during the hellish tearing of our Shrines ///

THANK GOD WE HAVE NUKES!!! /// 840 am (5/2)

sittin with slash, feedin ducks oats, smokin fat heart attack blunt -

5/2. c 4/4

from street salad blend: 1/5 ~ tobacco from blaine corner restaurant bar / 2/5 - menthol premiere first choice / 1/5 tobacco from streets following bellingham waterways / paths) ε, ν 1/5 marijuana pre-game / prior to horrid act-tearing [the real "government point" was actually not this, but the one after (4/30 night) right before frosty denounced it and left... Great Sunset Great Company, as this morning... This is to ensure the bits of marijuana are not "the Government" AKA act-tearing as d smoked this terpe blend just after the Hospital ///

5/2. c 4/4

c 3/4

5/1 ~~4/30~~ c 4/4

# Kids and Wheeled Sports Safety

**Every hour, nearly 50 children** visit emergency departments with an injury related to **bikes, scooters, skates or skateboards.**

**Serious head injuries (concussions, internal injuries and fractures)** made up 11% of ED visits across the four wheeled sports.

**Fractures** to the **shoulder, arm, elbow, wrist or hand** were the most frequent diagnoses for hospital admissions.

19% of hospital admissions for **scooter injuries** to children in 2015 were because of a **head injury.**




**Almost 40% of parents** of children ages 5-14 years indicated that their **child did not always wear a helmet** when participating in one of the four wheeled sports.

Among parents who say they **always wear a helmet** when riding a bike, 86% say their child also does. However, among parents who say they **never wear a helmet**, only 38% say their child always does.







**Reasons why** parents report their **children do not always wear a helmet:**

**47%** Parents think area is safe/View child as experienced/Don't see helmets as necessary.

**27%** Child finds helmet uncomfortable.

**24%** Other kids don't wear one.

**22%** Child thinks helmets aren't cool.

## Top Tips to Keep Kids Safe on Wheels

- All riders should wear a properly-fitted helmet. It is the best way to prevent head injuries and death.
- Ensuring correct fit of a helmet can increase comfort and use.
- Knee pads and elbow pads are recommended for scooters, skaters and skateboarders. Wrist guards are also recommended for skaters and skateboarders.

SAFE KIDS WORLDWIDE™